

Wentworth  
North

MacK's Cafe  
↓

Eric Dahlman

James  
Muller

Chase  
Duffy  
Cunningham  
= Dave Lick  
Jord Ann

## UTAH INFLUENZA IMMUNIZATION PROJECT

### INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at

Wm. W. W. W.

Wm. W. W.

Wm. W. W.

## UTAH INFLUENZA IMMUNIZATION PROJECT

### INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at

W. I.

that

Johnston

Kuhni-Midway

Port Johnson

Ally Green

Yelp Gault

(Raggs One)

Ally Gault +

Song Robert

## UTAH INFLUENZA IMMUNIZATION PROJECT

### INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at

Spauldine Ave  
New

Derek E. E. E. E. E.  
Mother's name  
(1) Ring with  
father's name  
Killed on the

Thank Kerby  
Derek E. E. E. E. E.  
New

## UTAH INFLUENZA IMMUNIZATION PROJECT

### INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at